

Three-body hadronic structure of low-lying $1/2^+$ Σ and Λ resonances

A. Martínez Torres, K. P. Khemchandani and E. Oset

Departamento de Física Teórica and IFIC, Centro Mixto Universidad de Valencia-CSIC,
Institutos de Investigación de Paterna, Aptd. 22085, 46071 Valencia, Spain.

Abstract

We discuss the dynamical generation of some low-lying $1/2^+$ Σ 's and Λ 's in two-meson one-baryon systems. These systems have been constructed by adding a pion in S -wave to the $\bar{K}N$ pair and its coupled channels, where the $1/2^-$ $\Lambda(1405)$ -resonance gets dynamically generated. We solve Faddeev equations in the coupled-channel approach to calculate the T -matrix for these systems as a function of the total energy and the invariant mass of one of the meson-baryon pairs. This squared T -matrix shows peaks at the energies very close to the masses of the strangeness -1, $1/2^+$ resonances listed in the particle data book.

1 Introduction

Several claims of excitation of a narrow resonance at ~ 1540 MeV were made in the pK_s final state in different experiments [1], which were performed to confirm the existence of the pentaquark state, Θ^+ [2]. The pK_s system could possess a total strangeness +1 or -1 and, hence, these states could very well correspond to, for example, a Σ^* [3, 4]. Actually, the existing Σ and Λ resonances [5], in this energy region, are not well understood. The poor status of these low-lying, $S = -1$ states is evident from the following facts: a) The spin-parity assignment for many of these states is unknown, e.g., for $\Sigma(1480)$, $\Sigma(1560)$, etc. b) the partial-wave analyses and production experiments have been often kept separately in the particle data group listings, *e.g.*, for $\Sigma(1620)$, $\Sigma(1670)$, c) other times, *e.g.*, in case of $\Lambda(1600)$, it is stated that existence of two resonances, in this energy region, is quite possible [5].

There are hints for some of them, like the $\Lambda(1600)$ and the $\Sigma(1660)$, to decay to three-body final states like $\pi^0\pi^0\Lambda$ and $\pi^0\pi^0\Sigma$, etc., [6, 7]. This indicates that the information of similar three-body contributions to the wave functions of the states, in this energy region, could be important and that this, so far unknown, information could lead to a better understanding of

the low-lying Σ and Λ resonances. The π - Σ interaction has been studied extensively in chiral unitary dynamics, where the $\Lambda(1405)$ gets dynamically generated (with a two pole structure [8]). Is it possible that adding another pion to this strongly attractive system could give rise to a three-body bound state? We have carried out a study of such systems for total spin-parity $(J^P) = 1/2^+$ and, indeed, find evidence for four Σ and two Λ known resonances [5] to have a three body, i.e., two-meson one-baryon, structure, which we discuss in the following text. The chiral dynamics has been used earlier in the context of the three-nucleon problems, *e.g.*, in [9]. This is the first time when the chiral dynamics is applied to solve the Faddeev equations for two-meson one-baryon systems.

2 Formalism

We solve Faddeev equations in the coupled channel approach. These coupled systems have been constructed by pairing up all the possible pseudoscalar mesons and $1/2^+$ baryons, which couple to strangeness $=-1$ and by adding a pion to the pair finally. We end up with twenty-two coupled channels for a fixed total charge [10]. In order to generate dynamically the $\Lambda(1405) S_{01}$ ($J^P = 1/2^-$) in the meson-baryon sub-systems, all the interactions have been written in the S -wave, which implies that the total J^P of the three body system is $1/2^+$.

The solution of the Faddeev equations,

$$T = T^1 + T^2 + T^3, \quad (1)$$

where the T^i partitions are written in terms of two body t -matrices (t^i) and three body propagators ($g = [E - H]^{-1}$) as

$$T^i = t^i + t^i g [T^j + T^k] \quad (i \neq j, i, j \neq k = 1, 2, 3), \quad (2)$$

requires off-shell two body t -matrices as an input. However, we found that the off-shell contributions of the two body t -matrices give rise to three-body forces which, when summed up for different diagrams, cancel the one arising directly from the chiral Lagrangian in the $SU(3)$ limit and for the case of small momentum transfer of the baryon [10]. In the realistic case, this sum was found to be only 5% of the total on-shell contribution [10]. Thus, it is reasonably accurate to study the problem by solving the Faddeev equations with the on-shell two-body t -matrices and by neglecting the three-body forces.

In this way, a term at second order in t in the Faddeev partitions (fig. 1), for instance,

$$t^1 g^{13} t^3, \quad (3)$$

is written as a product of the on-shell two body t -matrices t^1 and t^3 and the propagator g^{13} given as

$$g^{13} = \frac{1}{2E_2} \frac{1}{\sqrt{s} - E_1(\vec{k}'_1) - E_2(\vec{k}'_1 + \vec{k}_3) - E_3(\vec{k}_3) + i\epsilon}.$$




Figure 1: *The diagrammatic representation of the $t^1 g t^3$ term.*

Adding another interaction to the diagram in fig. 1 (see fig. 2), the expression $t^1 g^{13} t^3$ gets extended to




Figure 2: *The diagrammatic representation of $t^2 g^{21} t^1 g^{13} t^3$.*

$t^2 g^{21} t^1 g^{13} t^3$, which can be written explicitly as

$$t^2(s_{13}) \left[\int \frac{d\vec{k}''}{(2\pi)^3} \frac{1}{2E_1(\vec{k}'')} \frac{1}{2E_2(\vec{k}'' + \vec{k}_3)} \frac{2M_3}{2E_3(\vec{k}'' + \vec{k}'_2)} \right. \\ \times \frac{t^1(s_{23}(\vec{k}''))}{\sqrt{s} - E_1(\vec{k}'') - E_2(\vec{k}'_2) - E_3(\vec{k}'' + \vec{k}'_2) + i\epsilon} \\ \left. \times \frac{1}{\sqrt{s} - E_1(\vec{k}'') - E_2(\vec{k}'' + \vec{k}_3) - E_3(\vec{k}_3) + i\epsilon} \right] t^3(s_{12}),$$

where $s_{13} = (P - K'_2)^2$, $s_{12} = (P - K_3)^2$, $s_{23}(\vec{k}'') = (P - K'')^2$ with P, K_3, K'_2 and K'' representing the four momenta corresponding to the $\vec{P} = 0, \vec{k}_3, \vec{k}'_2$ and $\vec{k}''$ respectively. Our aim is to extract $t^1 g^{13} t^3$ out of the integral, which could simplify the calculations. The $t^2(s_{13})$ and $t^3(s_{12})$, in the equation above, depend on on-shell variables and can be factorized out of the loop integral but not t^1 and g^{13} [10]. This can be done if we re-arrange the loop integral as

$$t^2(s_{13}) \int \frac{d\vec{k}''}{(2\pi)^3} \frac{1}{2E_1(\vec{k}'')} \frac{2M_3}{2E_3(\vec{k}'' + \vec{k}'_2)} \frac{1}{\sqrt{s} - E_1(\vec{k}'') - E_2(\vec{k}'_2) - E_3(\vec{k}'' + \vec{k}'_2) + i\epsilon} \\ \times \left\{ t^1(s_{23}(\vec{k}'')) \frac{\frac{1}{2E_2(\vec{k}'' + \vec{k}_3)} \frac{1}{\sqrt{s} - E_1(\vec{k}'') - E_2(\vec{k}'' + \vec{k}_3) - E_3(\vec{k}_3) + i\epsilon}}{\frac{1}{2E_2(\vec{k}'_1 + \vec{k}_3)} \frac{1}{\sqrt{s} - E_1(\vec{k}'_1) - E_2(\vec{k}'_1 + \vec{k}_3) - E_3(\vec{k}_3) + i\epsilon}} \times [t^1(s_{23})]^{-1} \right\}$$

$$\times \left(t^1(s_{23}) \frac{1}{2E_2(\vec{k}'_1 + \vec{k}_3)} \frac{1}{\sqrt{s} - E_1(\vec{k}'_1) - E_2(\vec{k}'_1 + \vec{k}_3) - E_3(\vec{k}_3) + i\epsilon} t^3(s_{12}) \right).$$

The expression in the circular brackets, in the above equation, corresponds to $t^1 g^{13} t^3$ term which can be now extracted out of the loop integral since it depends on the on-shell variables. The off-shell dependence of t^1 and g^{13} has been included in the terms enclosed in the curly brackets, which we define as F^{213} and name as off-shell factor (which should not be confused with the off-shell “form-factor”, often used in the literature. The off-shell factor, here, is just the nomenclature for a mathematical expression which retains the off-shellness of the terms which are extracted out of the integral). The integral is now left with two propagators and the off-shell factor F^{213} , which together we re-express and define as a loop function, G^{213} ,

$$G^{213} = \int \frac{d\vec{k}''}{(2\pi)^3} \frac{1}{2E_1(\vec{k}'')}\frac{2M_3}{2E_3(\vec{k}'')}\frac{F^{213}(\sqrt{s}, \vec{k}'')}{\sqrt{s_{13}} - E_1(\vec{k}'') - E_3(\vec{k}'') + i\epsilon}$$

The diagram in fig. 2 can, hence, be re-written as $t^2 G^{213} t^1 g^{13} t^3$. The formalism has been developed following the above procedure, i.e., by replacing g by G , everytime a new interaction is added. This leads to another form of the Faddeev partitions (eq. (2)), which, after removing the terms corresponding to the disconnected diagrams and by denoting the rest of the equation as T_R , can be re-written as

$$T_R^{ij} = t^i g^{ij} t^j + t^i G^{ijk} T_R^{jk} + t^i G^{iji} T_R^{ji}, \quad i \neq j, j \neq k = 1, 2, 3. \quad (4)$$

The T_R^{ij} can be related to the Faddeev partitions (eq. (2)) as $T^i = t^i + T_R^{ij} + T_R^{ik}$, hence, giving six coupled equations instead of three (eq. (2)). These T_R^{ij} partitions correspond to the sum of all the possible diagrams with the last two interactions written in terms of t^i and t^j . This re-grouping of diagrams is done for the sake of convenience due to the different forms of the G^{ijk} functions. We define T_R as

$$T_R = \sum_{i \neq j=1}^3 T_R^{ij}, \quad (5)$$

which can be related to the sum of the Faddeev partitions (eq. (1)) as

$$T = t^1 + t^2 + t^3 + T_R. \quad (6)$$

3 Results and discussion

We shall construct the three-body T_R -matrices using the isospin symmetry, for which we must take an average mass for the isospin multiplets π (π^+ , π^0 , π^-), $\bar{K}$ ($\bar{K}^0$, K^-), K (K^+ , K^0), N (p , n), Σ (Σ^+ , Σ^0 , Σ^-) and Ξ (Ξ^0 , Ξ^-). In order to identify the nature of the resulting

states, we project the T_R -matrix on the isospin base. One appropriate base is the one where the states are classified by the total isospin of the three particles, I , and the total isospin of the two mesons, I_π in the case of two pions. Using the phase convention $|\pi^+\rangle = -|1, 1\rangle$, $|\pi^-\rangle = -|1, -1\rangle$, $|\Sigma^+\rangle = -|1, 1\rangle$ and $|\Xi^-\rangle = -|1/2, -1/2\rangle$ we have, for example, for the $\pi\pi\Sigma$ channel

$$\begin{aligned} |\pi^0 \pi^0 \Sigma^0\rangle &= |1, 0\rangle_\pi \otimes |1, 0\rangle_\pi \otimes |1, 0\rangle_\Sigma \\ &= \left\{ \sqrt{\frac{2}{3}} |2, 0\rangle - \sqrt{\frac{1}{3}} |0, 0\rangle \right\}_{\pi\pi} \otimes |1, 0\rangle_\Sigma \\ &= \sqrt{\frac{2}{5}} |I=3, I_\pi=2\rangle - \frac{2}{\sqrt{15}} |I=1, I_\pi=2\rangle - \\ &\quad - \sqrt{\frac{1}{3}} |I=1, I_\pi=0\rangle, \end{aligned}$$

where, I and I_π denote the total isospin of the three body system and that of the two pion system, respectively. Similarly, we write the other states in the isospin base.

After projecting the T_R -matrix (eq. (5)) on the isospin base, we square them and plot them as a function of the total energy ($\sqrt{s}$) of the two meson one baryon system and the invariant mass of particles 2 and 3 ($\sqrt{s_{23}}$). However, this choice is arbitrary, we could as well plot the squared amplitude, for example, as a function of $\sqrt{s_{12}}$ and $\sqrt{s_{13}}$.

In fig. 3, we show our results for $\pi\pi\Lambda$ in the $|I, I_\pi\rangle = |0, 0\rangle$ case, where a peak at 1740 MeV is clearly seen. The full width at half maximum of this peak is ~ 20 MeV. We identify this resonant structure with the $\Lambda(1810)$ in the particle data book [5]. It should be noted that, even though the status of this resonance in [5] is three-star, the associated pole positions vary from about 1750 MeV to 1850 MeV and the corresponding widths from 50-250 MeV.

We find evidence for another resonance in the isospin zero sector in the $\Lambda(1600)$ region. Apart from this, we find evidence for four low-lying Σ resonances: 1) for the $\Sigma(1560)$, thus predicting $J^P = 1/2^+$ for it, 2) the $\Sigma(1620)$, for which the partial-wave analyses and experimental findings are kept separately in the *PDG*, 3) the well-established $\Sigma(1660)$ and 4) the one-star $\Sigma(1770)$. All these results are summarized in the table 1, along with the states listed by the Particle Data Group.

References

- [1] DIANA Collaboration (V. V. Barmin *et al.*) , Phys. Atom. Nucl. **66**, 1715 (2003); HERMES Collaboration (A. Airapetian *et al.*) , Phys. Lett. B **585**, 213 (2004); SVD Collaboration (A. Aleev *et al.*) , Phys. Atom. Nucl. **68**, 974 (2005); ZEUS Collaboration (S. Chekanov *et al.*) , Phys. Lett. B **591**, 7 (2004); COSY-TOF Collaboration (M. Abdel-Bary *et al.*) , Phys. Lett. B **595**, 127 (2004), etc.
- [2] T. Nakano *et al.*, Phys. Rev. Lett. **91**, (2003) 012002.

Table 1: A comparison of the resonances found in this work with the states in *PDG*.

	Γ (<i>PDG</i>) (MeV)	Peak position (this work, MeV)	Γ (this work) (MeV)
Isospin = 1			
$\Sigma(1560)$	10 - 100	1590	70
$\Sigma(1620)$	10 - 100	1630	39
$\Sigma(1660)$	40 - 200	1656	30
$\Sigma(1770)$	60 - 100	1790	24
Isospin = 0			
$\Lambda(1600)$	50 - 250	1568	60
$\Lambda(1810)$	50 - 250	1740	20

- [3] Q. Zhao and F. E. Close, J. Phys. G **31**, L1 (2005).
- [4] F. E. Close and Q. Zhao, Phys. Lett. B **590**, 176 (2004).
- [5] W.-M. Yao *et al.*, J. Phys. G **33**, 1 (2006).
- [6] S. Prakhov *et al.*, Phys. Rev. C **70**, 034605 (2004).
- [7] S. Prakhov *et al.*, Phys. Rev. C **69** (2004) 042202.
S. Prakhov *et al.* [Crystall Ball Collaboration], Phys. Rev. C **70**, 034605 (2004).
- [8] D. Jido *et. al.*, Nucl. Phys. A **725**, 181 (2003).
- [9] E. Epelbaum *et. al.*, Phys. Rev. Lett. **86**, 4787 (2001).
- [10] A. Martinez Torres, K. P. Khemchandani and E. Oset, Phys. Rev. C **77**, 042203 (2008)




Figure 3: The $\Lambda(1810)$ resonance depicted in the squared amplitude for $\pi\pi\Lambda$ in the $|I, I_\pi\rangle = |0, 0\rangle$ configuration.